

Prescription Treatment[®] brand Alpine[®]

Cockroach Gel Bait Piston Can

KILLS: Cockroaches

FOR USE IN AND AROUND: Commercial, Institutional and Warehousing Establishments (including computer facilities, food/feed processing plants, hospitals, hotels, laboratories, motels, nursing homes, pet shops, restaurants, schools, supermarkets, warehouses and zoos), Residences and Transportation Equipment (aircraft, buses, boats/ships and trains)

ACTIVE INGREDIENT:

Dinotefuran, N-methyl-N'-nitro-N-[(tetrahydro-3-furanyl)methyl]guanidine 0.5%

OTHER INGREDIENTS:

99.5%

TOTAL: 100.0%

EPA Reg. No. 499-510

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED: Call a poison control center or doctor for treatment advice. Do not induce vomiting unless told to do so by the poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-832-HELP for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting. Do not use this product where a possibility of a shock hazard exists.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

READ ENTIRE LABEL. Use strictly in accordance with Precautionary Statements and Directions and with applicable state and federal regulations. Apply bait only in areas inaccessible to children and pets.

GENERAL INFORMATION

USE RESTRICTIONS

- Keep out of sunlight or excess heat.
- Do not apply bait to surfaces which may reach excessive temperatures. Examples include, but are not limited to: portions of stoves, ovens, grills, fume hoods, steam tables, toasters, fryers, dishwashers and hot water pipes.

SPOT TREATMENT

To apply gel, insert provided applicator straw into actuator. Apply gel in a 3 - 6 mm diameter spot (0.25 - 0.50 g). Approximately 10 spots per 100 ft² are recommended for use under conditions with low to moderate infestation. For more severe infestations, approximately 20 spots per 100 ft² are recommended. Apply many spots in areas of roach traffic and suspected harborage. Numerous small application spots are preferable to fewer larger application spots. Do not treat food preparation surfaces. Do not apply gel to areas that have been recently sprayed with insecticide. Do not spray insecticide products over gel placements as this may make gel repellent and reduce its effectiveness. Do not place treatments in locations where routine cleaning might remove bait or transfer bait to food or food preparation surfaces. This product will stick to surfaces that are not dusty or greasy. Reapply if gel placements are no longer visible and/or reinfestation occurs. Use paper towel to remove unwanted bait placements and then discard in trash.

CRACK & CREVICE[®] TREATMENT

This product can be applied as spots or as a line along cracks and crevices. Gel bait should be applied directly into cracks and crevices where insects hide. Place the applicator straw directly into cracks and crevices and deposit gel in small spots or in a continuous bead. Bait should be in the void area and not on open or exposed surfaces.

FOOD HANDLING ESTABLISHMENTS

Application within food areas of food handling establishments is limited to cracks and crevices where insects hide. To apply gel, insert provided applicator straw into actuator. Apply gel in a 3 - 6 mm diameter bead (0.25 - 0.50 g). Approximately 10 bait placements per 100 ft² are recommended for use under conditions with low to moderate infestation. For more severe infestations, approximately 20 bait placements per 100 ft² are recommended. Numerous small applications are preferable to fewer larger applications. Bait should not be used in open areas or on surfaces where contamination of food or food handling equipment may occur. Reapply gel if previous applications are no longer visible and/or reinfestation occurs.

APPLICATIONS IN AIRCRAFT

Application within passenger areas of aircraft is limited to Crack & Crevice treatment only. Apply this product in areas between elements of construction (such as areas under sinks, openings around pipes, under shelving, where shelving meets other shelves or the wall, area between and around microwaves, where support braces attach to the main structure and where seats are attached to the floor). Apply 10 - 20 bait spots per 100 ft² as described above. Do not apply bait to areas where food, utensils or food preparation surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food preparation and food holding surfaces as bait may be removed by washing. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface. During follow-up visits, inspect bait placements and reapply when previous applications are no longer visible and/or reinfestation occurs. Do not apply any bait in cockpit area.

OUTDOOR USE

Use for control of cockroaches. Inject into cracks and crevices around windows and doors, porches, screens, eaves, patios, garages, under stairways and other areas adjacent to structures where cockroaches may find harborage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry place away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site, according to label directions, or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not puncture or incinerate! Do not reuse or refill this container. Empty container by using the product according to label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly filled, contact your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured for:
Whitmire Micro-Gen Research Laboratories, Inc.[®]
by BASF Corporation
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
Questions? Call 1-800-777-8570
© 2010 Whitmire Micro-Gen Research Laboratories, Inc.[®]

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.