

- AND OTHER LISTED INSECTS
- FOR CONTROL OF TICKS WHICH MAY CARRY LYME DISEASE
  - ALSO FOR TREATMENT OF CLOTHING, TENTS, CAMPING GEAR AND: MOSQUITO NETTING, NETS, OUTSIDE OF SLEEPING BAGS, BACKPACKS AND GROUND CLOTHS EPA Reg. No. 45385-99-56

**KEEP OUT OF** REACH OF CHILDREN CAUTION

See side panels for additional Precautionary Statements.

ACTIVE INGREDIENT: 645-53-1]... OTHER INGREDIENTS:

TOTAL 100.00%

EPA Est. No. 88357-OH-01

Made in the U.S.A. © 2011

## **NET CONTENTS:**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Wash thoroughly after handling and before smoking or eating.

## FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person

 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. IF IN EYES

IF ON SKIN OR CLOTHING

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. IF INHALED

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

HOT LINE • You may also call 1-800-664-9042 for emergency medical treatment advice.

### **ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic organisms\_including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gotters or surface waters. Applying this product in calm weather when rainfs not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds.

Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

### PHYSICAL AND CHEMICAL HAZARDS

equipment due to the possibility of shock hazard. Do not allow product to freeze. Do not use this product in or on electrical

### STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area out of reach of

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the container be used, carefully following label directions and precautions.

CONTAINER DISPOSAL: IF EMPTY: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

IF PARTLY FILLED- Call your local waste agency or 1-800-253-2687 for disposal instructions. Never place unused product down any indoor or

SPILL: In case of spill, absorb liquid with absorbent material such as soil or clay, etc. sweep up and dispose of material according to PRODUCT DISPOSAL above.

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. Do not use on humans. Vacate room after treatment and ventitate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

Application is prohibited directly into sewers or drains or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during

or after application.

Ready-to-use and apply for most uses with the attached sprayer.

Manufactured for: J.T. EATON & CO., INC. 1393 East Highland Road Twinsburg, OH 44087 • U.S.A.• www.iteaton.com

### **Directions For Use continued**

For travelers who want to protect their luggage from picking up of bedbugs, spiders, cockroaches and other listed insects, spray the outer and inner surfaces of luggage with a light spray mist of this product. Allow the luggage to completely dry before use.

### INDOOR APPLICATION

INDOOR APPLICATION
Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensits should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Cover items that might water stain. Nonfood/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended. Do not apply to classrooms while in use.

### SURFACE SPRAYING:

TO KILL AND CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, ANTS, SILVERISH, SPIDERS, CRICKETS:
Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices, and infested areas of shelving.

TO CONTROL BED BUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floor. Repeat treatment as needed. Allow all treated articles to thoroughly dry before use.

TO CONTROL FLEAS (ADULT AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE) Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Allow to dry thoroughly before allowing nets to contact treated areas. before allowing pets to contact treated areas.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sun light, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all treated articles to dry thoroughly before use.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly

Close all doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths and Tobacco Moths: Direct spray at an upward angle. distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

INDOOR FOGGING: To kill exposed or accessible stages of the above named insects, use a rate of 1 quart to 1 gallon for 10,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Thoroughly ventilate before reoccupying.

### FOR USE ON DOGS:

FOR USE ON DOGS:

To Kill and control Fleas, Ticks, and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with pentials. Use protective gloves or mitts when applying product. Repeat application every two weeks, if necessary. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Do not use on dogs under 12 weeks of age. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathey our pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

### DO NOT USE ON CATS OR KITTENS.

### OUTDOOR APPLICATION

Do not water the treated area to the point of run-off. Do not make applications during rain. All outdoor applications must be limited to spot or crack -and-crevice treatments only, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawns, turf and other vegetation; 3) Applications to building foundations, up to a maximum height of 3 feet. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.

and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.

TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):
Use JT EATON KILLS BEDBUGS, TICKS AND MOSQUITOES as a residual treatment to control Ticks including the Deer Tick (Ixodes dammini), Lone Star Tick (Amblyomma americanum), Brown Dog Tick (Phipicephalus sanguinous) American Dog Tick (Demacentovariabilis) and Western Black-legged Tick (Ixodes pacificus). For each area listed below, apply until surface is wet (1/2 to 1 gallon for 750-1,000 sq. ft.)
Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter. Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

# CAMPING EQUIPMENT, MOSQUITO NETTING, NETS, TENTS, outsides of SLEEPING BAGS, BACKPACKS AND

CAMPING EQUIPMENT, MOSQUITO NETTING, NETS, TENTS, outsides of SLEEPING BAGS, BACKPACKS AND GROUND CLOTHS

For residual protection against ticks, chiggers and mosquitoes. Treat camping equipment as follows: This product should not be applied to clothing while it is being worn. Under NO Circumstances should bare skin or clothing on the body be treated. Do not spray directly on or otherwise allow it to come in contact with skin, face or eyes. This product is to be used as a clothing and gear treatment only. In case of accidental spraying of skin, face or eyes, see "FIRST AID".

Make all applications outdoors.

Make all applications outdoors.

For six week protection against ticks, chiggers and mosquitoes, select an outdoor area protected from the wind. Spray outer surfaces of clothing only (while NOT being worn) with a slow sweeping motion to lightly moisten the surface of the fabric, holding arm at a distance of 6 to 8 inches. (A complete outfit consists of shirt, trousers, and socks). Treat outside of each item of clothing for a minimum of 30 seconds on each side and allow 2 hours (4 hours under humid conditions) to dry prior to being worn. Pay particular attention to socks, trouser cuffs, and shirt cuffs. Treatment should moisten the surface of the fabric enough to cause a slight color change. Use approximately 2 to 3 ounces of this container to treat the outsides of one complete set of clothing. Use remainder on gear. Follow normal laundering procedures weekly. Reapply after six weeks and sixth laundering

and sixth laundering.

In heavily infested areas protection from ticks and chiggers will be improved by either wearing the cuffs inside footwear or taping pant cuffs around legs. Protection from mosquitoes can be further enhanced by applying an EPA approved repellent as directed, to face, hands and other exposed areas of the body.

Do not treat inside of sleeping bag. Spray exterior surfaces of tent only. Select a well ventilated outdoor area protected from wind and lay out gear to be treated. Hold container upright about 6 inches to 8 inches from surface of the article and spray with a slow sweeping motion to lightly moisten the surface of the fabric. Continue spraying over entire article, until the outer surface of the article is moist enough to cause a slight color change (darkening). Turn the article over and treat the other side. Hang treated gear and allow to dry for at least 2 hours (4 hours under humid conditions) before using. Do not retreat gear more than once every six weeks.

### NOTICE

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

**ORDER CODE 209-W**