





Prescription Treatment® brand PERMA-DUST®

Pressurized Boric Acid Dust

KILLS:

Ants, Booklice, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Centipedes, Chocolate Moths, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestides, Drugstore Beetles, Earwigs, Elm Leaf Beetles, Firebrats, Flour Beetles, Grain Weevils, Ground Beetles, Indianmeal Moths, Millipedes, Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders and Trogodermas.

FOR USE IN AND AROUND:

Apartments, Food Processing Plants, Food Storage Areas, Homes, Hospitals, Hotels, Inedible Product Areas of Meat Packing Plants, Motels, Nursing Homes, Restaurants and Other Food Handling Establishments, Schools*, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Planes**) and Warehouses and Other Commercial and Industrial Buildings.

- *Do not apply to classrooms when in use.
- **Do not use in aircraft cabins.

ACTIVE INGREDIENT:

 Boric Acid
 35.509

 OTHER INGREDIENTS:
 64.509

Contains petroleum distillates

TOTAL: 100.00%

EPA Reg. No. 499-384 KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment information.

NOTE TO PHYSICIAN: Contains petroleum distillates - vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Apply this product only as specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

SHAKE WELL BEFORE USING DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

All residue must be out of reach of children or domestic animals. Avoid contamination of food and foodstuffs. Do not use in aircraft cabins.

Before use, shake until agitator ball rattles. For best results, <u>use in upright position only</u>. Do not spray on plastic, painted or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, telephones, etc. where the powdery residue may foul or short electric contacts and circuits.

HOW TO USE WHITMIRE MICRO-GEN CRACK & CREVICE® INJECTION SYSTEM: Use Perma-Dust with the supplied actuator and injection tubes. Inject Perma-Dust into cracks and crevices or void spaces



where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Release approximately 1 second of product. Move injector tip along cracks while treating at the rate of 3 linear feet per second. For closed voids, calculate the void's cubic area and treat at the rate of 3 - 5 seconds per 3 cubic feet. For closed void areas, drill small access holes. Several holes may be required in long-running voids. This product may leave a white residue. This residue can normally be removed with a damp cloth or sponge.

INDOOR TREATMENTS

CENTIPEDES, COCKROACHES, CRICKETS, DER-MESTIDS, EARWIGS, MILLIPEDES, SCORPIONS, SOWBUGS AND SPIDERS: Inject with the prescribed dosage into cracks and crevices and void spaces where insects may be harboring. These may include, but are not limited to, openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewers, floor drains and meter boxes. BOOKLICE, FIREBRATS & SILVERFISH: Inject into cracks and crevices associated with baseboards, door frames, bookcases and other harborage sites.

ANTS: Inject cracks, crevices and voids where ants may be travelling and/or harboring, including, but not limited to, around doors, window frames, wall voids and other structural cracks and crevices and void areas.

STORED PRODUCT PESTS (Adult and Larval Stages) - Chocolate Moths, Drugstore Beetles, Flour Beetles, Grain Weevils, Indianmeal Moths, Trogodermas: Inject into cracks and crevices and/or voids where these insects may be harboring. Infested areas may include, but are not limited to, cracks and crevices in cabinets, pantries, wall voids and in machinery.

CARPENTER BEES: Locate bee entrance holes. Inject 5 - 10 seconds of product into entrance holes. Seal entrances after application.

APPLICATIONS IN FOOD HANDLING ESTABLISHMENTS

Food handling establishments are places other than private residences in which food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK & CREVICE TREATMENT ONLY: Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) are also considered food areas.

Apply directly into cracks and crevices and void areas using injection tube in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK & CREVICE TREATMENT, ARE NOT PERMITTED.

INDOOR AND OUTDOOR TREATMENTS

CARPENTER ANTS: Locate source of infestation or areas of suspected activity. Inject 5 - 10 seconds of product into insect tunnels and cavities. Treat points of entry or exits from voids such as around doors, windows and siding.

BOXELDER BUGS, CLOVER MITES, CLUSTER FLIES AND ELM LEAF BEETLES: Inject cracks and crevices and/or voids where these insects may be harboring or hibernating. Infested areas may include, but are not limited to, cracks and crevices, behind siding, wall voids, around window and door frames.

OUTDOOR TREATMENTS

ANTS, CENTIPEDES, CLOVER MITES, COCKROACH-ES, CRICKETS, EARWIGS, GROUND BEETLES, MIL-LIPEDES, PILLBUGS, SCORPIONS, SOWBUGS AND SPIDERS: Treat cracks around doors, windows, under siding and voids at foundation level, beneath guards on loading docks, into cracks in foundations and floors and in other places where these pests may enter building.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area away from heat or open flame and inaccessible to children.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product according to the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly filled, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc. 3568 Tree Court Industrial Blvd. St. Louis MO 63122-6682 © 2006 Whitmire Micro-Gen Research Laboratories, Inc.